



Test Definition: TTF40

TTF41 (SPT24) + p40 Immunostain, Technical Component Only

Overview

Useful For

Thyroid transcription factor 1 aids in the classification of carcinomas of unknown origin

p40 aids in the classification of carcinomas and lymphomas

Reflex Tests

| Test Id | Reporting Name | Available Separately | Always Performed |
|---------|--------------------------|----------------------|------------------|
| IHMTO | IHC Multiplex, Tech Only | No | No |

Testing Algorithm

[For the initial technical component only immunohistochemical \(IHC\) stain performed, the appropriate bill-only test ID will be reflexed and charged \(IHTOI\). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged \(IHTOA\).](#)

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain; **no pathologist interpretation is provided.**

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled**. Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

Shipping Instructions

Attach the green "Attention Pathology" address label (T498) and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Specimen Type: Tissue

Supplies: Immunostain Technical Only Envelope (T693)

Container/Tube: Immunostain Technical Only Envelope

Preferred:

-Formalin-fixed, paraffin-embedded tissue block

OR

-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Acceptable: None

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/pathology/digital-imaging/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/pathology/digital-imaging/#section3>

Forms

If not ordering electronically, complete, print, and send an [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

| | |
|---|--------|
| Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides Snowcoat slides | Reject |
|---|--------|

Specimen Stability Information

| Specimen Type | Temperature | Time | Special Container |
|---------------|---------------------|------|-------------------|
| TECHONLY | Ambient (preferred) | | |
| | Refrigerated | | |

Clinical & Interpretive

Clinical Information

Thyroid transcription factor 1 (TTF1) is a nuclear protein (detected by the chromogen 3,3'-diaminobenzidine) expressed in thyroid follicular cells, type II pneumocytes, and a subset of bronchial cells. The p40 antibody recognizes the deltaNp63 isoform of p63 (detected by the chromogen fast red). The predominant localization of p40 is in the basal layer of the stratified squamous and transitional epithelia. Given the relative specificity of TTF1 for cells of thyroid or lung origin, TTF1 is often included in a panel to identify the primary site for carcinomas of unknown origin. The p40 antibody may help to distinguish squamous cell carcinomas from other non-small cell carcinomas.

Reference Values

[This is not an orderable test. Order PATHC / Pathology Consultation. The consultant will determine the need for special stains.](#)

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity. If a control tissue is not included on the slide, a scanned image of the relevant quality control tissue is available upon request; call 855-516-8404.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- Minimize time slides are stored after being unpackaged
- Limit exposure to high humidity and heat
- Minimize exposure to plastics

Clinical Reference

1. Pelosi G, Fabbri A, Tamborini E, et al. Challenging lung carcinoma with coexistent deltaNp63/p40 and thyroid transcription factor-1 labeling within the same individual tumor cells. *J Thorac Oncol.* 2015;10(10):1500-1502
2. Whithaus K, Fukuoka J, Prihoda TJ, Jagirdar J. Evaluation of napsin A, cytokeratin 5/6, p63, and thyroid transcription factor 1 in adenocarcinoma versus squamous cell carcinoma of the lung. *Arch Pathol Lab Med.* 2012;136(2):155-162
3. Sterlacci W, Savic S, Schmid T, et al. Tissue-sparing application of the newly proposed IASLC/ATS/ERS classification of adenocarcinoma of the lung shows practical diagnostic and prognostic impact. *Am J Clin Pathol.* 2012;137(6):946-956

4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An introduction to the performance of immunohistochemistry. *Methods Mol Biol.* 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance

Method Description

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88344-TC

LOINC® Information

| Test ID | Test Order Name | Order LOINC® Value |
|---------|-----------------|--------------------|
|---------|-----------------|--------------------|

Test Definition: TTF40

TTF41 (SPT24) + p40 Immunostain, Technical
Component Only

| | | |
|-------|-----------------------------------|----------------------|
| TTF40 | TTF1 (SPT24) + p40 IHC, Tech Only | Order only;no result |
|-------|-----------------------------------|----------------------|

| Result ID | Test Result Name | Result LOINC® Value |
|-----------|-----------------------------------|----------------------|
| 602648 | TTF1 (SPT24) + p40 IHC, Tech Only | Bill only; no result |