



Test Definition: STICK

Tick-Borne Antibodies, Modified 2-Tier, ELISA,
Serum

Overview

Useful For

Diagnosis of Lyme disease and later stages of anaplasmosis, ehrlichiosis, and babesiosis

Sero-epidemiological surveys of the prevalence of the infection in certain populations

Profile Information

Test Id	Reporting Name	Available Separately	Always Performed
ANAP	Anaplasma phagocytophilum Ab, IgG,S	Yes	Yes
EHRC	Ehrlichia Chaffeensis (HME) Ab, IgG	Yes	Yes
BABG	Babesia microti IgG Ab, S	Yes	Yes
SLYME	Lyme Ab Modified 2-Tier w/Reflex, S	Yes	Yes

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
TLYME	Lyme IgM/IgG, WCS, EIA, S	Yes	No

Testing Algorithm

If the Lyme antibody result is positive or equivocal, then Lyme disease antibody confirmation will be performed at an additional charge.

For more information see [Acute Tickborne Disease Testing Algorithm](#).

Special Instructions

- [Acute Tickborne Disease Testing Algorithm](#)

Method Name

ANAP, EHRC, BABG: Immunofluorescence Assay (IFA)

SLYME: Enzyme-Linked Immunosorbent Assay (ELISA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Ordering Guidance

During the acute phase of an *Anaplasma phagocytophilum*, *Ehrlichia chaffeensis*, or *Babesia* infection, serologic tests are often nonreactive and detection by molecular assays is recommended. Polymerase chain reaction (PCR) testing is available to aid in the diagnosis of these cases; see EPCRB / *Ehrlichia/Anaplasma*, Molecular Detection, PCR, Blood and BABPB / *Babesia* species, Molecular Detection, PCR, Blood.

Specimen Required

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 1.35 mL Serum

Collection Instructions: Centrifuge and aliquot serum into a plastic vial.

Forms

If not ordering electronically, complete, print, and send [Infectious Disease Serology Test Request](#) (T916) with the specimen.

Specimen Minimum Volume

Serum: 1.1 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject
Heat-inactivated specimen	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	10 days	
	Frozen	14 days	

Clinical & Interpretive

Clinical Information

In North America, ticks are the primary vectors of infectious diseases.⁽¹⁾ Worldwide, ticks rank second only to mosquitoes in disease transmission. In the United States, tick-borne diseases include Lyme disease, Rocky Mountain spotted fever, human monocytic and granulocytic ehrlichiosis, babesiosis, tularemia, relapsing fever, and Colorado tick fever.

Symptoms of the various tick-vectoring diseases range from mild to life-threatening and significantly overlap. Early symptoms, which include fever, aches, and malaise, do not aid in distinguishing the various diseases. Because early treatment can minimize or eliminate the risk of severe disease, early detection is essential, yet patients may not have developed distinctive symptoms to help in the differential diagnosis. A tick-borne panel can assist in identifying the pathogen, allowing treatment to be initiated.

For information on specific diseases, see the individual test IDs.

Reference Values

ANAPLASMA PHAGOCYTOPHILUM ANTIBODY, IGG

<1:64

Reference values apply to all ages.

EHRlichia CHAFFEENSIS (HME) ANTIBODY, IGG

<1:64

Reference values apply to all ages.

BABESIA MICROTI IGG ANTIBODIES

<1:64

Reference values apply to all ages.

LYME ANTIBODY

Negative

Reference values apply to all ages.

Interpretation

Anaplasma phagocytophilum, *Ehrlichia chaffeensis*, and *Babesia microti*:

A positive immunofluorescence assay (titer > or =1:64) suggests current or previous infection. Seroconversion or a four-fold rise in titer is indicate active infection.

Lyme disease:

Negative:

Negative for antibodies to *Borrelia (Borrelia)* species causing Lyme disease. Negative results may occur in patients recently infected (< or =14 days) with *Borrelia burgdorferi*. If recent infection is suspected, repeat testing on a new sample collected in 7 to 14 days is recommended.

Equivocal or positive:

Not diagnostic. Supplemental testing in accordance with the modified two-tiered testing algorithm for Lyme disease has been ordered by reflex.

Cautions

Serology for IgG may be negative during the acute phase of infection (<7 days post-symptom onset), during which time detection using targeted nucleic acid amplification testing (eg, polymerase chain reaction: PCR) is recommended for *Anaplasma*, *Ehrlichia*, and *Babesia* infections

Detectable IgG-class antibodies typically appear within 7 to 10 days of post-symptom onset.

IgG-class antibodies may remain detectable months to years following prior infection. Therefore, a single time point-positive titer needs to be interpreted alongside other findings to differentiate recent versus past infection.

Other members of the *Ehrlichia* genus (eg, *Ehrlichia ewingii*) may not be detected by this assay.

Lyme disease:

A negative result does not exclude the possibility of infection with Lyme disease causing *Borrelia* species. Patients in the initial stages of Lyme disease and those who have been treated with antibiotics may not exhibit detectable antibody titers. Patients with clinical history, signs, or symptoms suggestive of Lyme disease should be retested in 2 to 4 weeks in the event that the initial test result is negative.

A positive result is not definitive evidence of infection with *Borrelia burgdorferi*. It is possible that other disease conditions may produce artifactual reactivity in the assay (eg, infectious mononucleosis, syphilis). All equivocal or positive results should be supplemented by immunoblot testing for IgM- and IgG-class antibodies in accordance with Centers for Disease Control and Prevention and the Association of State and Territorial Public Health Laboratory Directors (CDC/ASTPHLD) recommendations.

Patients infected with other members of the *B burgdorferi* sensu lato complex, including *Borrelia garinii*, *Borrelia afzelii*, and *Borrelia mayonii* will be detected by this assay; however, they cannot be differentiated.

This test should not be performed as a screening procedure for the general population. The predictive value of a positive or negative result depends on the prevalence of analyte (antibodies present to VlsE1 and pepC10 antigens) in a given population. Testing should only be performed when clinical evidence suggests the diagnosis of *Borrelia* infection or related etiological conditions observed by the physician.

Lyme serology should not be used for treatment monitoring, as IgG can remain for years post-resolution of infection. Instead, monitoring resolution of symptoms in response to treatment is recommended.

Clinical Reference

- Centers for Disease Control and Prevention (CDC). Tick-borne diseases of the United States: A Reference Manual for Health Care Providers. 5th ed. CDC; 2018. Available at www.cdc.gov/ticks/tickbornediseases/index.html
- Theel ES. The past, present and (possible) future of serologic testing for Lyme disease. J Clin Microbiol. 2016;54(5):1191-1196. doi:10.1128/JCM.03394-15

Performance

Method Description*Anaplasma phagocytophilum:*

The patient's serum is diluted and is placed in microscopic slide wells that have been coated with *Anaplasma phagocytophilum*-infected cells. After incubation, the slides are washed and a fluorescein-isothiocyanate conjugate is added to each well. The slides are then read using a fluorescence microscope and significant fluorescent staining of intracellular organisms constitutes a positive reaction. (Dumler JS, Asanovich KM, Bakken JS, Richter P, Kimsey R, Madigan JE. Serologic cross-reactions among *Ehrlichia equi*, *Ehrlichia phagocytophila*, and human granulocytic Ehrlichia. J Clin Microbiol. 1995;33[5]:1098-1103; Pancholi P, Kolbert CP, Mitchell PD, et al. *Ixodes dammini* as a potential vector of human granulocytic ehrlichiosis. J Infect Dis. 1995;172[4]:1007-1012; Dawson JE, Fishbein DB, Eng TR, Redus MA, Green NR. Diagnosis of human ehrlichiosis with the indirect fluorescent antibody test: kinetics and specificity. J Infect Dis. 1990;162[1]:91-95; package insert: Anaplasma phagocytophila IFA IgG. DiaSorin Molecular; 8/12/2016)

Ehrlichia chaffeensis:

The patient's serum is diluted and is placed in microscopic slide wells that have been coated with *Ehrlichia chaffeensis*-infected cells. After incubation, the slides are washed and a fluorescein-isothiocyanate conjugate is added to each well. The slides are then read using a fluorescence microscope and significant fluorescent staining of intracellular organisms constitutes a positive reaction. (Dumler JS, Asanovich KM, Bakken JS, Richter P, Kimsey R, Madigan JE. Serologic cross-reactions among *Ehrlichia equi*, *Ehrlichia phagocytophila*, and human granulocytic Ehrlichia. J Clin Microbiol. 1995;33[5]:1098-1103; Pancholi P, Kolbert CP, Mitchell PD, et al. *Ixodes dammini* as a potential vector of human granulocytic ehrlichiosis. J Infect Dis. 1995;172[4]:1007-1012; Dawson JE, Fishbein DB, Eng TR, Redus MA, Green NR. Diagnosis of human ehrlichiosis with the indirect fluorescent antibody test: kinetics and specificity. J Infect Dis. 1990;162[1]:91-95; package insert: Ehrlichia chaffeensis IFA IgG. DiaSorin Molecular; 8/12/2016)

Babesia microti:

This immunofluorescence assay detects antibodies against *Babesia microti*. The patient's serum is diluted and is placed in microscopic slide wells that have been coated with *B microti*-infected red blood cells from Syrian hamsters. After incubation, the slides are washed and a fluorescein-isothiocyanate conjugate is added to each well. The slides are then read using a fluorescence microscope and significant fluorescent staining of intraerythrocytic organisms constitutes a positive reaction. (Krause PJ, Telford SR 3rd, Ryan R, et al. Diagnosis of babesiosis: evaluation of a serologic test for the detection of *Babesia microti* antibody. J Infect Dis. 1994;169[4]:923-926; package insert: Babesia IFA IgG. DiaSorin Molecular; 8/12/2016)

Lyme disease:

The first-tier Lyme disease screening enzyme-linked immunosorbent assay (ELISA) used is the Zeus ELISA *Borrelia* VlsE1/pepC10 IgG/IgM test system. The Zeus ELISA *Borrelia* VlsE1/pepC10 IgG/IgM test system is designed to detect IgG- and IgM-class antibodies (not differentiated by the assay in the final result) in human sera to VlsE1 and pepC10 antigens. Diluted test sera are incubated in antigen coated microwells. Any antigen-specific antibody in the sample will bind to the immobilized antigen. The plate is washed to remove unbound antibody and other serum components. Peroxidase conjugated goat antihuman IgG and IgM are added to the wells and the plate is incubated. The conjugate will react with IgG and IgM antibodies immobilized on the plate. The wells are washed to remove unreacted conjugate. The microwells containing immobilized peroxidase conjugate are incubated with peroxidase substrate solution. Hydrolysis of the substrate by peroxidase produces a color change. After a period of time the reaction is stopped, and the color intensity of the solution is measured photometrically. The color intensity of the solution depends upon the antibody concentration in the original test sample. (Package insert: *Borrelia* VlsE1/pepC10 IgG/IgM Test System. Zeus Scientific,

Inc; Rev. Date 05/25/2021)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

2 to 4 days

Specimen Retention Time

14 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

See Individual Test IDs

CPT Code Information

- 86618
- 86666 x 2
- 86753
- 86617 x 2 (if appropriate)

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
STICK	Tick-Borne Abs w/ Lyme MTTA, S	103603-7

Result ID	Test Result Name	Result LOINC® Value
81157	Anaplasma phagocytophilum Ab, IgG,S	23877-4
81128	Babesia microti IgG Ab, S	16117-4
81478	Ehrlichia Chaffeensis (HME) Ab, IgG	47405-6

Test Definition: STICK

Tick-Borne Antibodies, Modified 2-Tier, ELISA,
Serum

SLYME	Lyme Ab Modified 2-Tier w/Reflex, S	83081-0
-------	-------------------------------------	---------