

## Overview

### Useful For

Aiding in the biochemical diagnosis of Krabbe disease using whole blood specimens

Follow-up of individuals affected with Krabbe disease

Follow-up testing after an abnormal newborn screening result for Krabbe disease

Monitoring of individuals at risk to develop late onset Krabbe disease

Monitoring of individuals with Krabbe disease after hematopoietic stem cell transplantation

### Genetics Test Information

Krabbe disease (globoid cell leukodystrophy) is an autosomal recessive lysosomal disorder caused by deficient activity of the enzyme galactocerebrosidase.

Krabbe disease is clinically variable; infantile-onset Krabbe disease is the most severe type, with rapid neurological regression resulting in early death.

### Testing Algorithm

If the patient has abnormal newborn screening result for Krabbe disease, immediate action should be taken. Refer to the appropriate American College of Medical Genetics and Genomics Newborn Screening ACT Sheet.(1,2)

The following are available:

[-Newborn Screen Follow-up for Krabbe Disease: Galactocerebrosidase](#)

[-Newborn Screen Follow-up for Krabbe Disease: Galactocerebrosidase and Psychosine](#)

### Special Instructions

- [Newborn Screen Follow-up for Krabbe Disease: Galactocerebrosidase](#)
- [Newborn Screen Follow-up for Krabbe Disease: Galactocerebrosidase and Psychosine](#)

### Highlights

Elevations in psychosine support a diagnosis of Krabbe disease; therefore, psychosine quantitation is a useful biomarker in determining if an individual has active disease. In addition, psychosine quantitation in red blood cells may be a valuable biomarker to monitor disease progression or treatment response in individuals of all ages.

Psychosine may also be elevated in saposin A cofactor deficiency, which results in a similar clinical phenotype to Krabbe disease, but patients typically have normal galactocerebrosidase activity in vitro.

### Method Name

Liquid Chromatography Tandem Mass Spectrometry (LC-MS/MS)

### NY State Available

---

Yes

## Specimen

### Specimen Type

Whole blood

### Ordering Guidance

This test is recommended for individuals of all ages, infancy through adulthood, for diagnosis or long-term monitoring of patients who have been treated or who are at risk of developing Krabbe disease.

If a result is needed within 24 hours, order PSY / Psychosine, Blood Spot.

### Shipping Instructions

1. Pre-analytical processing is performed Monday through Friday and Sunday. This test may be canceled if specimens are outside of stability when processing occurs. Collect and package specimens for arrival on days when processing is performed.

2. **Must be sent refrigerated.**

### Necessary Information

1. **Patient's age is required.**

2. Date of hematopoietic stem cell transplantation (HSCT), if performed.

### Specimen Required

#### Container/Tube:

**Preferred:** Lavender top (EDTA)

**Acceptable:** Green top (sodium heparin, lithium heparin)

**Specimen Volume:** 2 mL

**Collection Instructions:** Send whole blood specimen in original tube. **Do not aliquot.**

### Forms

[If not ordering electronically, complete, print, and send a Biochemical Genetics Test Request \(T798\)](#) with the specimen.

### Specimen Minimum Volume

0.5 mL

### Reject Due To

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Whole blood	Refrigerated	7 days	

---

## Clinical & Interpretive

### Clinical Information

Krabbe disease (globoid cell leukodystrophy) is an autosomal recessive lysosomal disorder caused by deficient activity of the enzyme galactocerebrosidase (GALC). GALC facilitates the lysosomal degradation of psychosine (galactosylsphingosine) and 3 other substrates, galactosylceramide, lactosylceramide, and lactosylsphingosine. Krabbe disease is caused by disease-causing variants in the *GALC* gene.

Severely affected individuals typically present between 3 to 6 months of age with increasing irritability and sensitivity to stimuli. Rapid neurodegeneration, including white matter disease, follows with death usually occurring by 2 years old. Some individuals have later onset forms of the disease that are characterized by ataxia, vision loss, weakness, and psychomotor regression presenting anywhere from 1 year to the seventh decade of life. The clinical course of Krabbe disease can be variable, even within the same family.

Newborn screening for Krabbe disease has been implemented in some states. The early (presymptomatic) identification and subsequent testing of infants at risk for Krabbe disease may be helpful in reducing the morbidity and mortality associated with this disease. While treatment is mostly supportive, hematopoietic stem cell transplantation has shown some success if performed prior to onset of neurologic damage.

Psychosine is 1 of 4 substrates degraded by GALC and is a neurotoxin at elevated concentrations. It has been shown to be elevated in patients with active Krabbe disease or with saposin A cofactor deficiency and, therefore, may be a useful biomarker for the presence of disease or disease progression.

Reduced or absent GALC in leukocytes (GALCW / Galactocerebrosidase, Leukocytes) or dried blood spots (PLSD / Lysosomal and Peroxisomal Disorders Screen, Blood Spot) along with elevated psychosine levels can indicate a diagnosis of Krabbe disease. Molecular sequencing of the *GALC* gene (GALC / Krabbe Disease, *GALC* Gene Sequencing with Deletion/Duplication, Varies) allows for detection of the disease-causing variants in affected patients and carrier detection in family members.

Individuals with a disease phenotype similar to Krabbe disease may have saposin A cofactor deficiency. Saposin A cofactor deficiency also results in elevated psychosine levels. Testing for this condition via molecular analysis of *PSAP* is useful in those with elevated psychosine and normal to moderately reduced GALC activity with normal molecular genetic *GALC* sequencing.

### Reference Values

Normal <5 pmol/g Hb

### Interpretation

An elevation of psychosine is indicative of Krabbe disease or saposin A cofactor deficiency.

### Cautions

Individuals with later onset disease, such as adult-onset Krabbe disease, may have a normal psychosine result.

This test is **not** capable of identifying carriers of Krabbe disease.

---

**Clinical Reference**

1. Newborn Screening ACT Sheet [Decreased galactocerebrosidase, elevated psychosine] Krabbe Disease (infantile form). American College of Medical Genetics and Genomics; 2021. Updated May 2022. Accessed October 2, 2025. Available at [www.acmg.net/PDFLibrary/Krabbe-Infantile.pdf](http://www.acmg.net/PDFLibrary/Krabbe-Infantile.pdf)
2. Newborn Screening ACT Sheet [Decreased galactocerebrosidase, mildly elevated psychosine] Krabbe Disease (late-onset form). American College of Medical Genetics and Genomics; 2021. Updated May 2022. Accessed October 2, 2025. Available [www.acmg.net/PDFLibrary/Krabbe-Later-Onset.pdf](http://www.acmg.net/PDFLibrary/Krabbe-Later-Onset.pdf)
3. Kwon JM, Matern D, Kurtzberg J, et al. Consensus guidelines for newborn screening, diagnosis and treatment of infantile Krabbe disease. *Orphanet J Rare Dis.* 2018;13(1):30. doi:10.1186/s13023-018-0766-x
4. Orsini JJ, Escolar ML, Wasserstein MP, et al. Krabbe disease. In: Adam MP, Mirzaa GM, Pagon R, eds. *GeneReviews* [Internet]. University of Washington, Seattle; 2000. Updated October 11, 2018. Accessed October 2, 2025. Available at [www.ncbi.nlm.nih.gov/books/NBK1238/](http://www.ncbi.nlm.nih.gov/books/NBK1238/)
5. Matern D, Basheeruddin K, Klug TL, et al. Newborn Screening for Krabbe Disease: Status Quo and Recommendations for Improvements. *Int J Neonatal Screen.* 2024;10(1):10. doi:10.3390/ijns10010010
6. Wenger DA, Escolar ML, Luzi P, Rafi MA. Krabbe disease (globoid cell leukodystrophy). In: Valle DL, Antonarakis S, Ballabio A, Beaudet AL, Mitchell GA, eds. *The Online Metabolic and Molecular Bases of Inherited Disease.* McGraw-Hill; 2019. Accessed October 2, 2025. Available at <https://ommbid.mhmedical.com/content.aspx?sectionid=225546481&bookid=2709>
7. Guenzel AJ, Turgeon CT, Nickander KK, et al. The critical role of psychosine in screening, diagnosis, and monitoring of Krabbe disease. *Genet Med.* 2020;22(6):1108-1118. doi:10.1038/s41436-020-0764-y
8. Thompson-Stone R, Ream MA, Gelb M, et al. Consensus recommendations for the classification and long-term follow up of infants who screen positive for Krabbe disease. *Mol Genet Metab.* 2021;134(1-2):53-59. doi:10.1016/j.ymgme.2021.03.016

**Performance****Method Description**

Psychosine is extracted from washed red blood cells and quantified using an isotopically labeled internal standard by liquid chromatography tandem mass spectrometry.(Unpublished Mayo method)

**PDF Report**

No

**Day(s) Performed**

Preanalytical processing: Monday through Friday, Sunday

Assay performed: Tuesday, Thursday

**Report Available**

3 to 7 days

**Specimen Retention Time**

Residual whole blood: 14 days; Lysate: 2 months

**Performing Laboratory Location**

---

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

82542

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
PSYR	Psychosine, RBC	93687-2

Result ID	Test Result Name	Result LOINC® Value
606152	Interpretation (PSYR)	59462-2
606145	Psychosine, RBC	93687-2
606151	Reviewed By	18771-6