



# Test Definition: 5BETH

Factor V Bethesda Units, Plasma

## Overview

### Useful For

Detecting and quantifying the presence and titer of a specific factor inhibitor directed against coagulation factor V

### Method Name

Only orderable as a reflex. For more information see:

- ALUPP / Lupus Anticoagulant Profile, Plasma
- ALBLD / Bleeding Diathesis Profile, Limited, Plasma
- ACBL / Bleeding Diathesis Profile, Comprehensive, Plasma
- APROL / Prolonged Clot Time Profile, Plasma
- 5INHE / Factor V Inhibitor Evaluation, Plasma

Optical Clot-Based

### NY State Available

Yes

## Specimen

### Specimen Type

Plasma Na Cit

### Ordering Guidance

If type of inhibitor is unknown, order APROL / Prolonged Clot Time Profile, Plasma.

### Specimen Required

Only orderable as a reflex. For more information see:

- ALUPP / Lupus Anticoagulant Profile, Plasma
- ALBLD / Bleeding Diathesis Profile, Limited, Plasma
- APROL / Prolonged Clot Time Profile, Plasma
- 5INHE / Factor V Inhibitor Evaluation, Plasma

### Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject
Gross icterus	Reject

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

## Clinical & Interpretive

### Clinical Information

Significant bleeding can result from the presence of a coagulation factor inhibitor and could be life threatening. Whether the inhibitor is present due to hemophilia or is of an acquired nature, it greatly complicates the treatment process of a decreased factor level. The titer of the inhibitor may determine the mode of treatment. Bethesda units are a standardization to give a uniform definition of an inhibitor.

### Reference Values

Only orderable as a reflex. For more information see:

- ALUPP / Lupus Anticoagulant Profile, Plasma
- ALBLD / Bleeding Diathesis Profile, Limited, Plasma
- ACBL / Bleeding Diathesis Profile, Comprehensive, Plasma
- APROL / Prolonged Clot Time Profile, Plasma
- 5INHE / Factor V Inhibitor Evaluation, Plasma

< or =0.5 Bethesda Units

### Interpretation

An interpretive report will be provided when testing is complete.

### Cautions

No significant cautionary statements

### Clinical Reference

1. Biggs R, Bidwell E. A method for the study of antihemophilic globulin inhibitors with reference to six cases. *Br J Haematol.* 1959;5:379-395
2. Hoyer LW: Factor VIII inhibitors. In: Hoyer LW, eds. *Progress in Clinical and Biological Research.* Vol 150. R Alan Liss Inc, 1984:87-98
3. Kasper CK, Aledort L, Aronson D, et al. Proceedings: A more uniform measurement of factor VIII inhibitors. *Thromb Diath Haemorrh.* 1975;34(2):61
4. Kasper C, Ewing N. Acquired inhibitors of plasma coagulation factors. *J Med Tech* 1986;38:431-439
5. Kottke-Marchant K, ed: *Laboratory Hematology Practice.* Wiley Blackwell Publishing; 2012
6. Hoffman R, Benz Jr EJ, Silberstein LE, et al, eds. *Hematology: Basic Principles and Practice.* 7th ed. Elsevier; 2018

## Performance

### Method Description

Heat inactivated patient plasma, undiluted and serially diluted, is mixed with an equal volume of normal pooled plasma (NPP). The NPP supplies the factor against which the inhibitor is directed in a known concentration. The patient plasma

mixtures, along with a control (Bethesda Pool) of diluted NPP are incubated at 37 degrees C for 2 hours, after which factor activity is measured. The factor activity in the undiluted patient and its serial dilutions are compared to the factor activity recovered in the Bethesda Pool. These values are then used to calculate Bethesda units. One Bethesda unit is defined as the amount of antibody that will destroy 50% of the coagulation factor activity in 2 hours. (Owen CA Jr, Bowie EJW, Thompson JH Jr. Diagnosis of Bleeding Disorders. 2nd ed. Little, Brown and Company; 1975; Meijer P, Verbruggen HW, Spannagi M. Clotting factors and inhibitors: Assays and interpretation. In: Kottke-Marchant K, ed. Laboratory Hematology Practice. Wiley Blackwell Publishing; 2012:435-446)

**PDF Report**

No

**Day(s) Performed**

Monday through Friday

**Report Available**

2 to 5 days

**Specimen Retention Time**

7 days

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

85335

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
5BETH	FV Bethesda Units, P	3191-4

Result ID	Test Result Name	Result LOINC® Value
607433	FV Bethesda Units, P	3191-4