



# Test Definition: CCMGG

Comprehensive Cardiomyopathy Gene Panel,  
Varies

## Overview

### Useful For

Providing a genetic evaluation for patients with a personal or family history suggestive of a hereditary form of cardiomyopathy

Establishing a diagnosis of a hereditary form of cardiomyopathy

### Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULFB	Fibroblast Culture for Genetic Test	Yes	No
CULAF	Amniotic Fluid Culture/Genetic Test	Yes	No
_STR1	Comp Analysis using STR (Bill only)	No, (Bill only)	No
_STR2	Add'l comp analysis w/STR (Bill Only)	No, (Bill only)	No
MATCC	Maternal Cell Contamination, B	Yes	No

### Genetics Test Information

This test utilizes next next-generation sequencing to detect single nucleotide and copy number variants in 83 genes associated with hereditary forms of cardiomyopathy: *ABCC9, ACAD9, ACADVL, ACTC1, ACTN2, AGL, ALMS1, ALPK3, BAG3, BRAF, CDH2, CPT2, CRYAB, CSRP3, DES, DMD, DNAJC19, DOLK, DSC2, DSG2, DSP, ELAC2, EMD, FHL1, FKRP, FKTN, FLNC, GAA, GLA, HCN4, HRAS, JPH2, JUP, KRAS, LAMP2, LMNA, LZTR1, MAP2K1, MAP2K2, MRAS, MTO1, MYBPC3, MYH7, MYL2, MYL3, MYLK3, MYPN, NEXN, NKX2-5, NRAS, PCCA, PCCB, PKP2, PLN, PPA2, PPCS, PRDM16, PRKAG2, PTPN11, RAF1, RBM20, RIT1, RYR2, SCN5A, SGCD, SHOC2, SLC22A5, SOS1, SOS2, TAZ (TAFAZZIN), TBX20, TCAP, TMEM43, TMEM70, TNNC1, TNNI3, TNNI3K, TNNT2, TPM1, TRIM63, TTN, TTR, and VCL*. See [Targeted Genes and Methodology Details for Comprehensive Cardiomyopathy Gene Panel](#) and Method Description for additional details.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, familial screening, and genetic counseling for hereditary forms of cardiomyopathy.

[Prior Authorization](#) is available for this assay.

### Testing Algorithm

For cord blood specimens that have an accompanying maternal blood specimen, maternal cell contamination studies will be performed at an additional charge.

### Special Instructions

- 
- [Informed Consent for Genetic Testing](#)
  - [Hereditary Cardiomyopathies and Arrhythmias: Patient Information](#)
  - [Informed Consent for Genetic Testing \(Spanish\)](#)
  - [Targeted Genes and Methodology Details for Comprehensive Cardiomyopathy Gene Panel](#)
  - [Comprehensive Cardiomyopathy Gene Panel \(CCMGG\) Prior Authorization Ordering Instructions](#)

**Method Name**

Sequence Capture and Targeted Next-Generation Sequencing followed by Polymerase Chain Reaction (PCR) and Sanger Sequencing

**NY State Available**

Yes

**Specimen****Specimen Type**

Varies

**Ordering Guidance**

Upon request and after initial testing is complete, WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies may be added to this test. To obtain more information about this option or add WESPR testing, call 800-533-1710.

Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To modify this panel via CGPH, please use the Cardiovascular/Connective Tissue/Dyslipidemia/Cerebrovascular/Primary Ciliary Dyskinesia disease state for step 1 on the [Custom Gene Ordering Tool](#).

Targeted testing for familial variants (also called site-specific or known mutations testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

**Necessary Information**

[Prior Authorization](#) is available, **but not required**, for this test. If proceeding with the prior authorization process, submit the required form with the specimen.

**Specimen Required**

**Patient Preparation:** A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. For information about testing patients who have received a hematopoietic stem cell transplant, call 800-533-1710.

**Submit only 1 of the following specimens:**

**Specimen Type:** Whole blood

---

**Container/Tube:** Lavender top (EDTA) or yellow top (ACD)

**Specimen Volume:** 3 mL

**Collection Instructions:**

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**
3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

**Additional Information:**

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.
3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers.

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:**

DNA Saliva Kit High Yield (T1007)

Saliva Swab Collection Kit (T786)

**Container/Tube:**

**Preferred:** High-yield DNA saliva kit

**Acceptable:** Saliva swab

**Specimen Volume:** 1 Tube if using T1007 or 2 swabs if using T786

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient (preferred) 30 days/Refrigerated 30 days

**Additional Information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

**Specimen Type:** Blood spot

**Supplies:** Card-Blood Spot Collection (Filter Paper) (T493)

**Container/Tube:**

**Preferred:** Collection card (Whatman Protein Saver 903 Paper)

**Acceptable:** PerkinElmer 226 filter paper or blood spot collection card

**Specimen Volume:** 2 to 5 Blood spots

**Collection Instructions:**

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see [How to Collect a Dried Blood Spot Sample](#).
2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
3. Do not expose specimen to heat or direct sunlight.

4. Do not stack wet specimens.
5. Keep specimen dry.

**Specimen Stability Information:** Ambient (preferred)/Refrigerated

**Additional Information:**

1. Blood spot specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from blood spots, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.
2. Due to lower concentration of DNA yielded from blood spot, it is possible that additional specimen may be required to complete testing.
3. For collection instructions, see [Blood Spot Collection Instructions](#)
4. For collection instructions in Spanish, see [Blood Spot Collection Card-Spanish Instructions](#) (T777)
5. For collection instructions in Chinese, see [Blood Spot Collection Card-Chinese Instructions](#) (T800)

**Specimen Type:** Extracted DNA

**Container/Tube:**

**Preferred:** Screw Cap Micro Tube, 2mL with skirted conical base

**Acceptable:** Matrix tube, 1mL

**Collection Instructions:**

1. The preferred volume is at least 100 µL at a concentration of 75 ng/µL.
2. Include concentration and volume on tube.

**Specimen Stability Information:** Frozen (preferred) 1 year/Ambient/Refrigerated

**Additional Information:** DNA must be extracted in a CLIA-certified laboratory or equivalent and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

**Forms**

1. **New York Clients-Informed consent is required.**

Document on the request form or electronic order that a copy is on file.

The following documents are available:

- [Informed Consent for Genetic Testing](#) (T576)
- [Informed Consent for Genetic Testing \(Spanish\)](#) (T826)
- 2. [Hereditary Cardiomyopathies and Arrhythmias Patient Information](#)
- 3. [Comprehensive Cardiomyopathy Panel \(CCMGG\) Prior Authorization Ordering Instructions](#)
- 4. [If not ordering electronically, complete, print, and send a Cardiovascular Test Request Form](#) (T724) with the specimen.

**Specimen Minimum Volume**

See Specimen Required

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

**Clinical & Interpretive****Clinical Information**

Cardiomyopathies are a group of disorders characterized by disease of the heart muscle. Cardiomyopathy can be caused by either inherited, genetic factors or nongenetic (acquired) causes, such as infection or trauma. When the presence or severity of the cardiomyopathy observed in a patient cannot be explained by acquired causes, genetic testing for the inherited forms of cardiomyopathy may be considered. Overall, cardiomyopathies are some of the most common genetic disorders. The inherited forms of cardiomyopathy include hypertrophic cardiomyopathy (HCM), dilated cardiomyopathy (DCM), arrhythmogenic cardiomyopathy (ACM), and left ventricular noncompaction (LVNC).(1)

The hereditary form of HCM is characterized by left ventricular hypertrophy in the absence of other cardiac or systemic causes that may cause hypertrophy of the heart muscle, such as longstanding, uncontrolled hypertension or aortic stenosis. The incidence of HCM in the general population is approximately 1:200 to 1:500, and it is estimated that 30% to 60% of cases can be attributed to a genetic etiology.(2) Hereditary forms of HCM are most often caused by genes encoding proteins of the cardiac sarcomere, the functional contractile unit of the heart muscle.

Hereditary forms of DCM are characterized by ventricular dilation with reduced cardiac performance in the absence of other cardiac or systemic causes that may cause dilation of the heart muscle, such as hypertension and ischemic heart disease. The incidence of DCM in the general population is approximately 1 in 2500, and it is estimated that approximately 50% of cases can be attributed to a genetic etiology.(3) Hereditary forms of DCM are most often caused by genes encoding proteins of the cardiac cytoskeleton and sarcomere.

Left ventricular noncompaction is characterized by prominent trabeculations of the left ventricle with trabecular recesses extending into the ventricular cavity. The incidence of LVNC in the general population is estimated to be 1 in 5000.(3) It is currently unclear if LVNC represents a genetically distinct form of cardiomyopathy, as many familial cases of LVNC have been linked to the same genes associated with other forms of hereditary cardiomyopathies and many affected individuals also meet diagnostic criteria for DCM or HCM.(3,4)

Arrhythmogenic cardiomyopathy is characterized by the presence of arrhythmogenic cardiac muscle in the absence of ischemic, hypertensive, or valvular cardiac disease. Arrhythmogenic right ventricular cardiomyopathy (ARVC), the most well-defined form of ACM, is characterized by the breakdown of the myocardium and replacement of right ventricular muscle tissue with fibrofatty tissue, resulting in an increased risk of arrhythmia and sudden death. In some cases, there may also be left ventricular involvement. The prevalence of ARVC (genetic and acquired) is estimated to be 1 in 2000 to 1 in 5000 in the general population.(5)

Hereditary forms of cardiomyopathy may be an isolated finding or may be a feature of an underlying systemic condition. Hereditary forms of cardiomyopathy can follow autosomal dominant, autosomal recessive, X-linked, and digenic

---

patterns of inheritance. Mitochondrial inheritance is also possible, however, genes associated with mitochondrial inheritance of cardiomyopathy are not assessed on this panel.

**Reference Values**

An interpretive report will be provided.

**Interpretation**

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.<sup>(6)</sup> Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

**Cautions**

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact the Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

This analysis targets single and multi-exon deletions/duplications; however, in some instances single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the genes included on the panel may occur. In these instances, genes included in the ordered test are provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. However, copy number variants for genes not listed in the Method Description are typically not reported

---

or interpreted for haploinsufficiency/triplosensitivity. CMACB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the genes included on the panel.

This test is not designed to detect low levels of mosaicism or to differentiate between somatic and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent non-leukocyte reduced blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time.

#### Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.<sup>(6)</sup> Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. Incidental findings may include but are not limited to, results related to the sex chromosomes. These findings will be carefully reviewed to determine whether they will be reported.

#### Clinical Reference

1. Hershberger RE, Givertz MM, Ho CY, et al. Genetic evaluation of cardiomyopathy—a heart failure society of America practice guideline. *J Card Fail.* 2018;24(5):281-302. doi:10.1016/j.cardfail.2018.03.004
2. Ommen SR, Ho CY, Asif IM, et al. 2024 AHA/ACC/AMSSM/HRS/PACES/SCMR Guideline for the Management of Hypertrophic Cardiomyopathy: A Report of the American Heart Association/American College of Cardiology Joint Committee on Clinical Practice Guidelines. *Circulation.* 2024;149(23):e1239-e1311. doi:10.1161/CIR.0000000000001250
3. Bozkurt B, Colvin M, Cook J, et al. Current diagnostic and treatment strategies for specific dilated cardiomyopathies: a

---

scientific statement from the American Heart Association [published correction appears in *Circulation*. 2016;134(23):e652]. *Circulation*. 2016;134(23):e579-e646. doi:10.1161/CIR.0000000000000455

4. Aung N, Doimo S, Ricci F, et al. Prognostic significance of left ventricular noncompaction: Systematic review and meta-analysis of observational studies. *Circ Cardiovasc Imaging*. 2020;13(1):e009712. doi:10.1161/CIRCIMAGING.119.009712

5. Corrado D, Zorzi A, Cipriani A, et al. Evolving Diagnostic Criteria for Arrhythmogenic Cardiomyopathy. *J Am Heart Assoc*. 2021;10(18):e021987. doi:10.1161/JAHA.121.021987

6. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med* 2015;17(5):405-424. doi:10.1038/gim.2015.30

## Performance

### Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing is performed to test for the presence of variants in coding regions, and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletion-insertions (delins) less than 40 base pairs (bp), and above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Comprehensive Cardiomyopathy Gene Panel](#) for details regarding the targeted genes analyzed for each test and specific gene regions not routinely covered. (Unpublished Mayo method)

Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

Genes analyzed: *ABCC9, ACAD9, ACADVL, ACTC1, ACTN2, AGL, ALMS1, ALPK3, BAG3, BRAF, CDH2, CPT2, CRYAB, CSRP3, DES, DMD, DNAJC19, DOLK, DSC2, DSG2, DSP, ELAC2, EMD, FHL1, FKR, FKTN, FLNC, GAA, GLA, HCN4, HRAS, JPH2, JUP, KRAS, LAMP2, LMNA, LZTR1, MAP2K1, MAP2K2, MRAS, MTO1, MYBPC3, MYH7, MYL2, MYL3, MYLK3, MYPN, NEXN, NKX2-5, NRAS, PCCA, PCCB, PKP2, PLN, PPA2, PPCS, PRDM16, PRKAG2, PTPN11, RAF1, RBM20, RIT1, RYR2, SCN5A, SGCD, SHOC2, SLC22A5, SOS1, SOS2, TAZ (TAFAZZIN), TBX20, TCAP, TMEM43, TMEM70, TNNC1, TNNI3, TNNI3K, TNNT2, TPM1, TRIM63, TTN, TTR, and VCL*

### PDF Report

Supplemental

### Day(s) Performed

---

Varies

**Report Available**

21 to 35 days

**Specimen Retention Time**

Whole blood: 25 days (if available); Saliva: 30 days (if available); Extracted DNA: 3 months, Blood Spots: 1 year (if available)

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

81439

**Prior Authorization**

Insurance preauthorization is available for this testing; forms are available.

Patient financial assistance may be available to those who qualify. Patients who receive a bill from Mayo Clinic Laboratories will receive information on eligibility and how to apply.

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
CCMGG	Comprehensive Cardiomyopathy Panel	107547-2

Result ID	Test Result Name	Result LOINC® Value
617184	Test Description	62364-5
617185	Specimen	31208-2
617186	Source	31208-2
617187	Result Summary	50397-9

## Test Definition: CCMGG

Comprehensive Cardiomyopathy Gene Panel,  
Varies

617188	Result	82939-0
617189	Interpretation	69047-9
617190	Additional Results	82939-0
617191	Resources	99622-3
617192	Additional Information	48767-8
617193	Method	85069-3
617194	Genes Analyzed	48018-6
617195	Disclaimer	62364-5
617196	Released By	18771-6