



# Test Definition: AUTOG

Autoinflammatory Disorders Gene Panel,  
Varies

## Overview

### Useful For

Providing a comprehensive genetic evaluation for patients with a personal or family history suggestive of an inherited autoinflammatory disorder

Establishing a diagnosis of a monogenic autoinflammatory disorder, allowing for appropriate management and surveillance for disease features based on the gene or variant involved

Identifying variants within genes known to be associated with monogenic autoinflammatory disorders, allowing for predictive testing of at-risk family members

### Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
CULAF	Amniotic Fluid Culture/Genetic Test	Yes	No
_STR1	Comp Analysis using STR (Bill only)	No, (Bill only)	No
_STR2	Add'l comp analysis w/STR (Bill Only)	No, (Bill only)	No
CULFB	Fibroblast Culture for Genetic Test	Yes	No
MATCC	Maternal Cell Contamination, B	Yes	No

### Genetics Test Information

This test utilizes next-generation sequencing to detect single nucleotide and copy number variants in 117 genes associated with autoinflammatory disorders: *ACP5, ADA, ADA2, ADAM17, ADAR, AIRE, ALPI, AP3B1, AP3D1, ARPC1B, ASAH1, C1QA, C1QB, C1QC, C1R, C1S, C2, CARD11, CARD14, CASP10, CASP8, CD3G, CD40LG, CD48, CD55, CDC42, COPA, CTLA4, DDX58 (RIGI), DNASE1, DNASE1L3, DNASE2, DOCK8, FADD, FOXP3, GATA2, HAVCR2, ICOS, IFIH1, IKBKG, IL10, IL10RA, IL10RB, IL1RN, IL2RA, IL2RB, IL2RG, IL36RN, ISG15, ITCH, ITGB2, ITK, JAK1, LACC1, LIG4, LPIN2, LRBA, LSM11, LYN, LYST, MEFV, MVK, NLRC4, NLRP1, NLRP12, NLRP3, NOD2, OAS1, OTULIN, PIK3CD, PIK3R1, PLCG2, POLA1, POMP, PRF1, PRKCD, PSMA3, PSMB10, PSMB4, PSMB8, PSMB9, PSMG2, PSTPIP1, RAB27A, RBCK1, RIPK1, RNASEH2A, RNASEH2B, RNASEH2C, RNF31, RNU7-1, SAMD9L, SAMHD1, SH2D1A, SH3BP2, SKIV2L (SKIC2), SLC29A3, SLC37A4, STAT1, STAT2, STAT3, STIM1, STING1, STX11, STXBP2, TLR7, TNFAIP3, TNFRSF1A, TPP2, TREX1, TRNT1, UNC13D, USP18, WAS, WDR1, XIAP, and ZAP70.* See [Targeted Genes and Methodology Details for Autoinflammatory Disorders Gene Panel](#) for details regarding the targeted gene regions evaluated by this test.

Identification of a disease-causing variant may assist with diagnosis, prognosis, clinical management, recurrence risk assessment, familial screening, and genetic counseling for a heritable form of autoinflammatory disease.

---

**Testing Algorithm****Skin biopsy:**

For skin biopsy or cultured fibroblast specimens, fibroblast culture will be performed at an additional charge. If viable cells are not obtained, the client will be notified.

**Cord blood:**

For cord blood specimens that have an accompanying maternal blood specimen, maternal cell contamination studies will be performed at an additional charge.

**Special Instructions**

- [Informed Consent for Genetic Testing](#)
- [Blood Spot Collection Card-Spanish Instructions](#)
- [Blood Spot Collection Card-Chinese Instructions](#)
- [Informed Consent for Genetic Testing \(Spanish\)](#)
- [Blood Spot Collection Instructions](#)
- [Inborn Errors of Immunity, Autoimmunity, and Autoinflammatory Disease Patient Information](#)
- [Targeted Genes and Methodology Details for Autoinflammatory Disorders Gene Panel](#)

**Method Name**

Sequence Capture and Amplicon-Based Next-Generation Sequencing (NGS)/Polymerase Chain Reaction (PCR)/Droplet Digital Polymerase Chain Reaction (ddPCR)/Quantitative Real-Time Polymerase Chain Reaction (qPCR)/Sanger Sequencing as needed

**NY State Available**

Yes

**Specimen****Specimen Type**

Varies

**Ordering Guidance**

Patients who have had a previous bone marrow transplant from an allogenic donor should not have testing performed on blood, bone marrow, or saliva because any results generated will reflect the genome of the donor rather than the recipient. Testing on patients who have an active hematologic malignancy or hematologic disorder with clonal proliferation may identify both somatic mutations and germline variants, which may result in test failure or necessitate follow-up testing to determine whether the detected variant is germline or somatic. For these patients, testing a skin biopsy or cultured fibroblasts is recommended. For instructions for testing patients who have received a bone marrow transplant or have an active hematologic disorder, call 800-533-1710. For more information see Cautions.

Upon request and after initial testing is complete, WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies may be added to this test. To obtain more information about this option or add WESPR testing, call 800-533-1710.

---

Customization of this panel and single gene analysis for any gene present on this panel are available. For more information see CGPH / Custom Gene Panel, Hereditary, Next-Generation Sequencing, Varies. To modify this panel via CGPH, use the Inborn Errors of Immunity/Bone Marrow Failure/Telomeropathy/Pulmonary Fibrosis/Very Early Onset IBD/Pancreatitis disease state for step 1 on the [Custom Gene Ordering Tool](#).

Targeted testing for familial variants (also called site-specific or known variants testing) is available for the genes on this panel. See FMTT / Familial Variant, Targeted Testing, Varies. To obtain more information about this testing option, call 800-533-1710.

### Additional Testing Requirements

**For cord blood specimens:** Maternal cell contamination (MCC) studies are available. **Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal specimens under separate order numbers.** Cord blood testing will proceed without MCC studies, but results may be compromised if MCC is present.

### Specimen Required

**Patient Preparation:** A previous hematopoietic stem cell transplant from an allogenic donor will interfere with testing. Call 800-533-1710 for instructions for testing patients who have received a hematopoietic stem cell transplant.

### Submit only 1 of the following specimens:

**Specimen Type:** Whole blood

**Container/Tube:** Lavender top (EDTA) or yellow top (ACD)

**Specimen Volume:** 3 mL

#### Collection Instructions:

1. Invert several times to mix blood.
2. Send whole blood specimen in original tube. **Do not aliquot.**
3. Whole blood collected postnatal from an umbilical cord is also acceptable. See Additional Information.

**Specimen Stability Information:** Ambient (preferred) 4 days/Refrigerated 4 days/Frozen 4 days

#### Additional Information:

1. Specimens are preferred to be received within 4 days of collection. Extraction will be attempted for specimens received after 4 days, and DNA yield will be evaluated to determine if testing may proceed.
2. To ensure minimum volume and concentration of DNA are met, the requested volume must be submitted. Testing may be canceled if DNA requirements are inadequate.
3. For postnatal umbilical cord whole blood specimens, maternal cell contamination studies are recommended to ensure test results reflect that of the patient tested. A maternal blood specimen is required to complete maternal cell contamination studies. Order MATCC / Maternal Cell Contamination, Molecular Analysis, Varies on both the cord blood and maternal blood specimens under separate order numbers.

**Specimen Type:** Skin biopsy

**Supplies:** Fibroblast Biopsy Transport Media (T115)

**Container/Tube:** Sterile container with any standard cell culture media (eg, minimal essential media, RPMI 1640). The solution should be supplemented with 1% penicillin and streptomycin.

**Specimen Volume:** 4-mm Punch

---

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical or Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Cultured fibroblasts

**Source:** Skin

**Container/Tube:** T-25 flask

**Specimen Volume:** 2 Flasks

**Collection Instructions:** Submit confluent cultured fibroblast cells from a skin biopsy from another laboratory. Cultured cells from a prenatal specimen will not be accepted.

**Specimen Stability Information:** Ambient (preferred) <24 hours/Refrigerated <24 hours

**Additional Information:**

1. Specimens are preferred to be received within 24 hours of collection. Culture and extraction will be attempted for specimens received after 24 hours and will be evaluated to determine if testing may proceed.
2. A separate culture charge will be assessed under CULFB / Fibroblast Culture for Biochemical and Molecular Testing. An additional 3 to 4 weeks are required to culture fibroblasts before genetic testing can occur.

**Specimen Type:** Extracted DNA

**Container/Tube:**

**Preferred:** Screw Cap Micro Tube, 2 mL with skirted conical base

**Acceptable:** Matrix tube, 1 mL

**Collection Instructions:**

1. The preferred volume is at least 100 µL at a concentration of 75 ng/µL.
2. Include concentration and volume on tube.

**Specimen Stability Information:** Frozen (preferred) 1 year/Ambient/Refrigerated

**Additional Information:** DNA must be extracted in a CLIA-certified laboratory, or equivalent, and must be extracted from a specimen type listed as acceptable for this test (including applicable anticoagulants). Our laboratory has experience with Chemagic, Puregene, Autopure, MagnaPure, and EZ1 extraction platforms and cannot guarantee that all extraction methods are compatible with this test. If testing fails, one repeat will be attempted, and if unsuccessful, the test will be reported as failed and a charge will be applied. If applicable, specific gene regions that were unable to be interrogated due to DNA quality will be noted in the report.

**Specimen Type:** Blood spot

**Supplies:** Card-Blood Spot Collection (Filter Paper) (T493)

**Container/Tube:**

**Preferred:** Collection card (Whatman Protein Saver 903 Paper)

**Acceptable:** PerkinElmer 226 filter paper or blood spot collection card

**Specimen Volume:** 2 to 5 Blood spots

**Collection Instructions:**

1. An alternative blood collection option for a patient older than 1 year is a fingerstick. For detailed instructions, see [How to Collect a Dried Blood Spot Sample](#).

- 
2. Let blood dry on the filter paper at ambient temperature in a horizontal position for a minimum of 3 hours.
  3. Do not expose specimen to heat or direct sunlight.
  4. Do not stack wet specimens.
  5. Keep specimen dry

**Specimen Stability Information:** Ambient (preferred)/Refrigerated

**Additional Information:**

1. Blood spot specimens are acceptable but not recommended. Multiple extractions will be required to obtain sufficient yield for supplemental analysis, and there is significant risk for test failure due to insufficient DNA.
2. Due to lower concentration of DNA yielded from blood spot, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.
3. For collection instructions, see [Blood Spot Collection Instructions](#)
4. For collection instructions in Spanish, see [Blood Spot Collection Card-Spanish Instructions](#) (T777)
5. For collection instructions in Chinese, see [Blood Spot Collection Card-Chinese Instructions](#) (T800)

**Specimen Type:** Saliva

**Patient Preparation:** Patient should not eat, drink, smoke, or chew gum 30 minutes prior to collection.

**Supplies:**

DNA Saliva Kit High Yield (T1007)

Saliva Swab Collection Kit (T786)

**Container/Tube:**

**Preferred:** High-yield DNA saliva kit

**Acceptable:** Saliva swab

**Specimen Volume:** 1 Tube if using T1007 or 2 swabs if using T786

**Collection Instructions:** Collect and send specimen per kit instructions.

**Specimen Stability Information:** Ambient (preferred) 30 days/Refrigerated 30 days

**Additional Information:** Saliva specimens are acceptable but not recommended. Due to lower quantity/quality of DNA yielded from saliva, some aspects of the test may not perform as well as DNA extracted from a whole blood sample. When applicable, specific gene regions that were unable to be interrogated will be noted in the report. Alternatively, additional specimen may be required to complete testing.

**Forms**

1. **New York Clients-Informed consent is required.** Document on the request form or electronic order that a copy is on file. The following documents are available:

-[Informed Consent for Genetic Testing](#) (T576)

-[Informed Consent for Genetic Testing \(Spanish\)](#) (T826)

2. [Inborn Errors of Immunity, Autoimmunity and Autoinflammatory Disease Patient Information](#)

**Specimen Minimum Volume**

See Specimen Required

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Varies	Varies		

## Clinical & Interpretive

### Clinical Information

Systemic autoinflammatory disorders result from dysregulation of the innate immune system and are characterized by a hyperinflammatory state with elevated acute phase reactants. These disorders may present at any age, but symptoms often begin in childhood with unexplained fever that may be accompanied by a rash. While these features can mimic infections or hematological neoplasias, the inflammatory lesions are noncancerous and sterile. Additional features may be present and highly variable, depending on the organ or organs impacted by cytokine amplification loops and sterile inflammation. Symptoms may involve the gastrointestinal (GI) tract (eg, serositis, abdominal pain, early-onset inflammatory bowel disease), bone, eyes (eg, uveitis), musculoskeletal system (eg, arthritis and arthralgias), central nervous system (eg, meningitis), or other tissues. Some autoinflammatory disorders are also associated with an increased risk of developing amyloid A (AA) amyloidosis. These disorders include familial Mediterranean fever (FMF), tumor necrosis factor receptor-associated periodic syndrome (TRAPS), cryopyrin-associated autoinflammatory syndrome (CAPS), and hyper-IgD syndrome/mevalonate kinase deficiency (HIDS/MKD).

Autoinflammatory disorders are classified by molecular pathogenesis or clinical features. Pathophysiologic classification is based on the pathway or cytokine that drives disease, such as interleukin (IL)-1, interferon, nuclear factor kappa B, and IL-18. Disease classification based on clinical features often focuses on skin involvement or fever duration and frequency. Age of onset, triggers, and additional organ system involvement are also used to classify these disorders and aid clinical diagnosis.

The genetic basis of many heritable autoinflammatory disorders has been identified. Autoinflammatory disorders may be inherited in an autosomal recessive, autosomal dominant, or X-linked manner. Disease-causing variants may also arise *de novo*. The inheritance pattern appears more complicated for some disorders. For example, FMF is typically inherited in an autosomal recessive manner. However, some affected individuals appear to have only one disease-causing alteration. For other autoinflammatory disorders, cases of digenic and oligogenic inheritance have also been described. Inheritance may also be multifactorial, requiring an environmental component along with low-penetrance variants. One example is Yao syndrome, a recently described clinical entity characterized by recurrent fever, dermatitis, inflammatory arthritis, and GI symptoms in most affected individuals. While some variants in *NOD2* have been reported in association with Yao syndrome, they are relatively common among the general population. They may confer an increased risk for developing Yao syndrome but are not diagnostic and appear insufficient to cause disease by themselves. Some disorders such as PFAPA (periodic fever, aphthous stomatitis, pharyngitis, adenitis) syndrome, systemic juvenile idiopathic arthritis, adult-onset Still disease, and Behcet disease have significant phenotypic overlap with monogenic autoinflammatory conditions, but a genetic cause of these disorders has not been identified.

Finally, several examples of post-zygotic (mosaic or somatic) genetic alterations causing autoinflammatory disorders have been described. While it may be possible to identify mosaic variants, this test is primarily intended for the identification of germline variants and the diagnosis of inherited monogenic autoinflammatory disorders.

Determining the underlying genetic cause of an autoinflammatory condition may help guide treatment decisions. For

---

example, colchicine is an effective therapy for many patients with FMF, but some patients may not respond. Instead, these individuals, and others affected by a subset of autoinflammatory disorders, may respond to IL-1 blocking therapies. Anakinra, riloncept, and canakinumab are several examples of medications that target IL-1. However, another subset of autoinflammatory disorders is not responsive to IL-1 blockade, such as proteasome-associated autoinflammatory syndromes (PRAAS), CANDLE (chronic atypical neutrophilic dermatosis with lipodystrophy), deficiency of IL-36 receptor antagonist, and CAMPS (CARD14-mediated psoriasis). Medications that target other components of the IL-1 pathway are under development. In addition, medications that target other pathways (eg, anti-tumor necrosis factor, anti-IL-6, and JAK-inhibitors) have demonstrated efficacy in some patients.

**Reference Values**

An interpretive report will be provided.

**Interpretation**

All detected variants are evaluated according to American College of Medical Genetics and Genomics recommendations.<sup>(1)</sup> Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance.

**Cautions**

Clinical Correlations:

Test results should be interpreted in the context of clinical findings, family history, and other laboratory data. Misinterpretation of results may occur if the information provided is inaccurate or incomplete.

If testing was performed because of a clinically significant family history, it is often useful to first test an affected family member. Detection of a reportable variant in an affected family member would allow for more informative testing of at-risk individuals.

To discuss the availability of additional testing options or for assistance in the interpretation of these results, contact Mayo Clinic Laboratories genetic counselors at 800-533-1710.

Technical Limitations:

Next-generation sequencing may not detect all types of genomic variants. In rare cases, false-negative or false-positive results may occur. The depth of coverage may be variable for some target regions; assay performance below the minimum acceptable criteria or for failed regions will be noted. Given these limitations, negative results do not rule out the diagnosis of a genetic disorder. If a specific clinical disorder is suspected, evaluation by alternative methods can be considered.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. Confirmation of select reportable variants will be performed by alternate methodologies based on internal laboratory criteria.

This test is validated to detect 95% of deletions up to 75 base pairs (bp) and insertions up to 47 bp. Deletions-insertions (delins) of 40 or more bp, including mobile element insertions, may be less reliably detected than smaller delins.

Deletion/Duplication Analysis:

---

This analysis targets single and multi-exon deletions/duplications; however, in some instances, single exon resolution cannot be achieved due to isolated reduction in sequence coverage or inherent genomic complexity. Balanced structural rearrangements (such as translocations and inversions) may not be detected.

Deletion/duplication events that extend past the genes included on the panel may occur. In these instances, genes included in the ordered test are provided on the report and interpreted, and genomic breakpoints are reported if they are confirmed. However, copy number variants for genes not listed in the Method Description are typically not reported or interpreted for haploinsufficiency/triplosensitivity. CMCB / Chromosomal Microarray, Congenital, Blood; WESPR / Panel to Whole Exome Sequencing Reflex Test, Varies; or WGSDX / Whole Genome Sequencing for Hereditary Disorders, Varies is recommended for a full interpretation of deletions/duplications predicted to extend past the genes included on the panel.

This test is not designed to detect low levels of mosaicism or differentiate between somatic mutations and germline variants. If there is a possibility that any detected variant is somatic, additional testing may be necessary to clarify the significance of results.

Genes may be added or removed based on updated clinical relevance. For detailed information regarding gene specific performance and technical limitations, see Method Description or contact a laboratory genetic counselor.

If the patient has had an allogeneic hematopoietic stem cell transplant or a recent non-leukoreduced blood transfusion, results may be inaccurate due to the presence of donor DNA. Call Mayo Clinic Laboratories for instructions for testing patients who have received a bone marrow transplant.

#### Reclassification of Variants:

Currently, it is not standard practice for the laboratory to systematically review previously classified variants on a regular basis. The laboratory encourages healthcare professionals to contact the laboratory at any time to learn how the classification of a particular variant may have changed over time. Due to broadening genetic knowledge, it is possible that the laboratory may discover new information of relevance to the patient. Should that occur, the laboratory may issue an amended report.

#### Variant Evaluation:

Evaluation and categorization of variants are performed using published American College of Medical Genetics and Genomics and the Association for Molecular Pathology recommendations as a guideline.<sup>(1)</sup> Other gene-specific guidelines may also be considered. Variants are classified based on known, predicted, or possible pathogenicity and reported with interpretive comments detailing their potential or known significance. Variants classified as benign or likely benign are not reported.

Multiple in silico evaluation tools may be used to assist in the interpretation of these results. The accuracy of predictions made by in silico evaluation tools is highly dependent upon the data available for a given gene, and periodic updates to these tools may cause predictions to change over time. Results from in silico evaluation tools should be interpreted with caution and professional clinical judgment.

Rarely, incidental or secondary findings may implicate another predisposition or presence of active disease. These findings will be carefully reviewed to determine whether they will be reported.

### Clinical Reference

1. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424
2. Hashkes PJ, Laxer RM, Simon A. *Textbook of Autoinflammation*. Springer Nature; 2019
3. Tangye SG, Al-Herz W, Bousfiha A, et al. Human inborn errors of immunity: 2022 update on the classification from the International Union of Immunological Societies Expert Committee. *J Clin Immunol*. 2022;42(7):1473-1507
4. Rood JE, Behrens EM. Inherited autoinflammatory syndromes. *Annu Rev Pathol*. 2022;17:227-249
5. Gutierrez MJ, Lapidus SK. Systemic autoinflammatory diseases. *Rheum Dis Clin North Am*. 2022;48(1):371-395
6. Broderick L, Hoffman HM. IL-1 and autoinflammatory disease: biology, pathogenesis and therapeutic targeting. *Nat Rev Rheumatol*. 2022;18(8):448-463

### Performance

#### Method Description

Next-generation sequencing (NGS) and/or Sanger sequencing are performed to test for the presence of variants in coding regions and intron/exon boundaries of the genes analyzed, as well as some other regions that have known disease-causing variants. The human genome reference GRCh37/hg19 build was used for sequence read alignment. At least 99% of the bases are covered at a read depth over 30X. Sensitivity is estimated at above 99% for single nucleotide variants, above 94% for deletions/insertions (delins) less than 40 base pairs (bp), and above 95% for deletions up to 75 bp and insertions up to 47 bp. NGS and/or a polymerase chain reaction (PCR)-based quantitative method is performed to test for the presence of deletions and duplications in the genes analyzed. Supplemental PCR-based methods are used to detect a large deletion in *IKBKG* and an inversion in *UNC13D*. A supplemental droplet digital PCR method is used to detect deletions and duplications in *C1R*. Confirmation of select reportable variants may be performed by alternate methodologies based on internal laboratory criteria.

There may be regions of genes that cannot be effectively evaluated by sequencing or deletion and duplication analysis as a result of technical limitations of the assay, including regions of homology, high guanine-cytosine (GC) content, and repetitive sequences. See [Targeted Genes and Methodology Details for Autoinflammatory Disorders Gene Panel](#) for details regarding the targeted gene regions identified by this test.(Unpublished Mayo method)

#### Genes analyzed:

*ACP5, ADA, ADA2, ADAM17, ADAR, AIRE, ALPI, AP3B1, AP3D1, ARPC1B, ASAH1, C1QA, C1QB, C1QC, C1R, C1S, C2, CARD11, CARD14, CASP10, CASP8, CD3G, CD40LG, CD48, CD55, CDC42, COPA, CTLA4, DDX58 (RIGI), DNASE1, DNASE1L3, DNASE2, DOCK8, FADD, FOXP3, GATA2, HAVCR2, ICOS, IFIH1, IKBKG, IL10, IL10RA, IL10RB, IL1RN, IL2RA, IL2RB, IL2RG, IL36RN, ISG15, ITCH, ITGB2, ITK, JAK1, LACC1, LIG4, LPIN2, LRBA, LSM11, LYN, LYST, MEFV, MVK, NLRC4, NLRP1, NLRP12, NLRP3, NOD2, OAS1, OTULIN, PIK3CD, PIK3R1, PLCG2, POLA1, POMP, PRF1, PRKCD, PSMA3, PSMB10, PSMB4, PSMB8, PSMB9, PSMG2, PSTPIP1, RAB27A, RBCK1, RIPK1, RNASEH2A, RNASEH2B, RNASEH2C, RNF31, RNU7-1, SAMD9L, SAMHD1, SH2D1A, SH3BP2, SKIV2L (SKIC2), SLC29A3, SLC37A4, STAT1, STAT2, STAT3, STIM1, STING1, STX11, STXBP2, TLR7, TNFAIP3, TNFRSF1A, TPP2, TREX1, TRNT1, UNC13D, USP18, WAS, WDR1, XIAP, and ZAP70*

**PDF Report**

Supplemental

**Day(s) Performed**

Varies

**Report Available**

28 to 42 days

**Specimen Retention Time**

Whole blood 28 days (if available); Saliva: 30 days (if available); Extracted DNA: 3 months; Blood spots: 1 year (if available)

**Performing Laboratory Location**

Mayo Clinic Laboratories - Rochester Main Campus

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

**CPT Code Information**

81443

88233-Tissue culture, skin, solid tissue biopsy (if appropriate)

88240-Cryopreservation (if appropriate)

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
AUTOG	Autoinflammatory Gene Panel	107548-0

Result ID	Test Result Name	Result LOINC® Value
620093	Test Description	62364-5
620094	Specimen	31208-2
620095	Source	31208-2
620096	Result Summary	50397-9
620097	Result	82939-0

## Test Definition: AUTOG

Autoinflammatory Disorders Gene Panel,  
Varies

620098	Interpretation	69047-9
620099	Additional Results	82939-0
620100	Resources	99622-3
620101	Additional Information	48767-8
620102	Method	85069-3
620103	Genes Analyzed	82939-0
620104	Disclaimer	62364-5
620105	Released By	18771-6