



Test Definition: CD13

CD13 Immunostain, Technical Component
Only

Overview

Useful For

Marker of myeloid lineage

Reflex Tests

Test Id	Reporting Name	Available Separately	Always Performed
IHTOI	IHC Initial, Tech Only	No	No
IHTOA	IHC Additional, Tech Only	No	No

Testing Algorithm

For the initial technical component only immunohistochemical (IHC) stain performed, the appropriate bill-only test ID will be reflexed and charged (IHTOI). For each additional technical component only IHC stain performed, an additional bill-only test ID will be reflexed and charged (IHTOA).

Method Name

Immunohistochemistry (IHC)

NY State Available

Yes

Specimen

Specimen Type

TECHONLY

Ordering Guidance

This test includes only technical performance of the stain; **no pathologist interpretation is provided.**

Technical component only stains **should not** be ordered with PATHC / Pathology Consultation. If ordered with PATHC, the technical component stains **will be canceled**. Any immunohistochemistry (IHC)/in situ hybridization (ISH) stain performed as a part of the PATHC will be performed at the reviewing pathologist's discretion at an additional charge.

Shipping Instructions

Attach the green pathology address label and the pink Immunostain Technical Only label included in the kit to the outside of the transport container.

Specimen Required

Specimen Type: Tissue

Supplies: Immunostain Technical Only Envelope (T693)

Container/Tube: Immunostain Technical Only Envelope

Preferred:

-Formalin-fixed, paraffin-embedded tissue block

OR

-2 Unstained, positively charged glass slides (25- x 75- x 1-mm) per test ordered; sections 4-microns thick

Acceptable: None

Digital Image Access

1. Information on accessing digital images of immunohistochemical (IHC) stains and the manual requisition form can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/>
2. Clients ordering stains using a manual requisition form will not have access to digital images.
3. Clients wishing to access digital images must place the order for IHC stains electronically. Information regarding digital imaging can be accessed through this website: <https://news.mayocliniclabs.com/ihc-stains/#FAQ>

Forms

If not ordering electronically, complete, print, and send a [Immunohistochemical \(IHC\)/In Situ Hybridization \(ISH\) Stains Request](#) (T763) with the specimen.

Reject Due To

Wet/frozen tissue Cytology smears Nonformalin fixed tissue Nonparaffin embedded tissue Noncharged slides ProbeOn slides Snowcoat slides	Reject
---	--------

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
TECHONLY	Ambient (preferred)		
	Refrigerated		

Clinical & Interpretive

Clinical Information

CD13 (cluster of differentiation 13) plays roles in peptide metabolism (brush border membranes of small intestine, renal proximal tubules, and placenta), cell growth and differentiation, and phagocytosis. CD13 is normally expressed on myeloid lineage cells, including granulocytes and monocytes. It is also expressed on non-hematolymphoid cells including endothelial cells and fibroblasts and is present in a soluble form in plasma. This immunostain may be useful as a marker of myeloid lineage in acute leukemias.

Interpretation

This test does not include pathologist interpretation, only technical performance of the stain. If interpretation is required, order PATHC / Pathology Consultation for a full diagnostic evaluation or second opinion of the case.

The positive and negative controls are verified as showing appropriate immunoreactivity.

Interpretation of this test should be performed in the context of the patient's clinical history and other diagnostic tests by a qualified pathologist.

Cautions

Age of a cut paraffin section can affect immunoreactivity. Stability thresholds vary widely among published literature and are antigen dependent. Best practice is for paraffin sections to be cut within 6 weeks.

The charge of glass slides can be affected by environmental factors and subsequently may alter slide staining. Sending unsuitable glass slides can result in inconsistent staining due to poor slide surface chemistry.

Best practices for storage of positively charged slides:

- Minimize time slides are stored after being unpackaged
- Limit exposure to high humidity and heat
- Minimize exposure to plastics

Clinical Reference

1. Yamanaka C, Wada H, Eguchi H, et al. Clinical significance of CD13 and epithelial mesenchymal transition (EMT) markers in hepatocellular carcinoma. *Jpn J Clin Oncol*. 2018;48(1):52-60
2. Kessler T, Baumeier A, Brand C, et al. Aminopeptidase N (CD13): expression, prognostic impact, and use as therapeutic target for tissue factor induced tumor vascular infarction in soft tissue sarcoma. *Transl Oncol*. 2018;11(6):1271-1282
3. Dominguez JM, Perez-Chacon G, Guillen MJ, et al. CD13 as a new tumor target for antibody-drug conjugates: validation with the conjugate MI130110. *J Hematol Oncol*. 2020;13(1):32
4. Magaki S, Hojat SA, Wei B, So A, Yong WH. An Introduction to the Performance of Immunohistochemistry. *Methods Mol Biol*. 2019;1897:289-298. doi:10.1007/978-1-4939-8935-5_25

Performance**Method Description**

Immunohistochemistry on sections of paraffin-embedded tissue.(Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

Until staining is complete.

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been cleared, approved, or is exempt by the US Food and Drug Administration and is used per manufacturer's instructions. Performance characteristics were verified by Mayo Clinic in a manner consistent with CLIA requirements.

CPT Code Information

88342-TC, primary

88341-TC, if additional IHC

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
CD13	CD13 IHC, Tech Only	Order only;no result

Result ID	Test Result Name	Result LOINC® Value
70687	CD13 IHC, Tech Only	Bill only; no result