



# Test Definition: FDAIP

Diabetes Autoimmune Profile

## Overview

### Useful For

Diabetes autoantibodies assessment is helpful in identifying and managing patients at risk for development of type 1 diabetes.

### Method Name

Enzyme-Linked Immunosorbent Assay (ELISA), Radioimmunoprecipitation

### NY State Available

No

## Specimen

### Specimen Type

Serum

### Specimen Required

#### Collection Container/Tube:

**Preferred:** Serum gel

**Acceptable:** Red top

**Submission Container/tube:** Plastic vial

**Specimen Volume:** 2.5 mL

#### Collection Instructions:

1. Within 24 hours of collection, centrifuge and aliquot 2.5 mL of serum into a plastic vial.
2. Freeze within 24 hours and send frozen.

### Specimen Minimum Volume

1 mL (Note: This volume does **not** allow for repeat testing)

### Reject Due To

Gross hemolysis	Reject
Thawing	Cold OK; Warm reject
Lipemia	Reject
EDTA plasma	Reject

### Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
---------------	-------------	------	-------------------

Serum	Frozen	7 days	
-------	--------	--------	--

## Clinical & Interpretive

### Reference Values

Insulin Antibodies

<5.0 uU/mL: Negative

> or =5.0 uU/mL: Positive

Anti GAD 65 Antibodies

<5.0 U/mL: Negative

> or =5.0 U/mL: Positive

IA-2 Autoantibodies

<7.5 U/mL: Negative

> or =7.5 U/mL: Positive

ZNT8 Antibodies

<15 U/mL: Negative

> or =15 U/mL: Positive

Reference ranges apply to all ages.

### Interpretation

Published positivity rates for diabetes autoantibodies in new-onset type 1 diabetes patients listed below are based on the combined analysis of GAD-65, ICA 512, insulin antibodies, and ZNT8 antibodies. The combined analysis has a 98% autoimmunity detection rate, with 1.8% of type 1 diabetic individuals remaining as autoantibody-negative.(1)

Positive rate in new-onset type 1 diabetes patients:

GAD-65 antibodies = 68% positive

ICA 512 antibodies = 72% positive

Insulin antibodies = 55% positive

ZnT8 antibodies = 63% positive

An increase in the number of positive antibodies is associated with a higher likelihood of type 1 diabetes.

Less than 3% of type 2 diabetics have positive antibodies.

### Clinical Reference

1. Wenzlau JM, Juhl K, Yu L, et al. The cation efflux transporter ZnT8 (Slc30A8) is a major autoantigen in human type 1 diabetes. Proc Natl Acad Sci U S A. 2007;104(43):17040-17045. doi:10.1073/pnas.0705894104

## Performance

**PDF Report**

No

**Day(s) Performed**

Varies

**Report Available**

10 to 19 days

**Performing Laboratory Location**

Esoterix Endocrinology

**Fees & Codes****Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

**Test Classification**

Insulin Antibodies: This test was developed and its performance characteristics determined by LabCorp. It has not been cleared or approved by the Food and Drug Administration.

**CPT Code Information**

86337

86341x3

**LOINC® Information**

Test ID	Test Order Name	Order LOINC® Value
FDAIP	Diabetes Autoimmune Profile	Not Provided

Result ID	Test Result Name	Result LOINC® Value
Z6377	Insulin Antibodies	8072-1
Z6378	Anti GAD 65 Antibodies	56540-8
Z6387	IA-2 Autoantibodies	31209-0
Z6388	ZNT8 Antibodies	76651-9
Z6389	Type 1 Diabetes Interpretation	8251-1