



Test Definition: F9_IS

Factor IX Inhibitor Screen, Plasma

Overview

Useful For

Detecting the presence of a specific factor inhibitor directed against coagulation factor IX

Special Instructions

- [Coagulation Guidelines for Specimen Handling and Processing](#)

Method Name

Only orderable as a reflex. For more information see:

9INHE / Factor IX Inhibitor Evaluation, Plasma

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

APROL / Prolonged Clot Time Profile, Plasma

Optical Clot-Based

NY State Available

Yes

Specimen

Specimen Type

Plasma Na Cit

Specimen Required

Only orderable as a reflex. For more information see:

9INHE / Factor IX Inhibitor Evaluation, Plasma

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

APROL / Prolonged Clot Time Profile, Plasma

For more information see [Coagulation Guidelines for Specimen Handling and Processing](#).

Specimen Minimum Volume

Platelet poor plasma: 2 mL

Reject Due To

Gross hemolysis	Reject
-----------------	--------

Gross lipemia	Reject
Gross icterus	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Plasma Na Cit	Frozen	14 days	

Clinical & Interpretive**Clinical Information**

Patient plasma, normal pooled plasma (NPP), and a mixture of patient plasma and NPP are each tested for a specific factor, incubated at 37 degrees C for 1 hour, and then retested for the same factor. In addition, a new mixture of patient plasma and NPP is prepared using the incubated plasmas and tested after the 1-hour incubation. The percentage of the recovered factor for each individual plasma and mixture being tested is calculated and compared. The procedure demonstrates the effect of a specific coagulation factor inhibitor on that factor present in normal pooled plasma, over a specific period of time.

An inhibitor directed against a coagulation factor may arise due to multiple exposures from transfusions in a patient deficient in that factor (as in the case of persons with hemophilia), in response to certain diseases, or be drug induced. Nonspecific inhibitors may also be present in patients that will prolong screening tests (eg, prothrombin time and activated partial thromboplastin time). This test is used to qualitatively identify an inhibitor to a specific coagulation factor.

Reference Values

Only orderable as a reflex. For more information see:

9INHE / Factor IX Inhibitor Evaluation, Plasma

ALUPP / Lupus Anticoagulant Profile, Plasma

ALBLD / Bleeding Diathesis Profile, Limited, Plasma

APROL / Prolonged Clot Time Profile, Plasma

Negative

Interpretation

When testing is complete, if factor activity results fall within clinically normal ranges, an interpretive comment will be provided noting that inhibitor testing was not indicated and, therefore, not performed. If factor activity indicates the performance of inhibitor screen testing, an interpretive comment will be provided noting the presence or absence of a factor IX inhibitor.

Cautions

Occasionally, a potent lupus-like anticoagulant may cause false-positive testing for a specific factor inhibitor (eg, factor VIII or IX).

Clinical Reference

1. Bowie EJW, Thompson JH Jr, Didisheim P, Owen CA Jr. Mayo Clinic Laboratory Manual of Hemostasis. WB Saunders Company; 1971:111-115
2. Chitlur M, Warriar I, Rajpurkar M, et al. Inhibitors in factor IX deficiency a report of the ISTH-SSC international FIX inhibitor registry (1997-2006). Haemophilia. 2009;15(5):1027-1031
3. Kottke-Marchant K. Laboratory Hematology Practice. Wiley Blackwell Publishing; 2012
4. Favaloro EJ, Lippi G. Hemostasis and Thrombosis, Methods and Protocols. Humana Press; 2017

Performance

Method Description

The factor IX inhibitor screen is performed on the Instrumentation Laboratory ACL TOP. The assay consists of measuring the factor IX activity (prothrombin time based assay) before and after incubation of a mixture of normal plasma and patient's plasma for 1 hour at 37 degrees C. Interpretation of the presence or absence of the indication of a factor IX inhibitor is determined by comparing the factor IX activity results and the calculated expected values. (Owen CA Jr, Bowie EJW, Thompson JH Jr. The Diagnosis of Bleeding Disorders. 2nd ed. Little Brown and Company; 1975:14-144; Meijer P, Verbruggen HW, Spannagi M: Clotting factors and inhibitors: Assays and interpretation. In: Kottke-Marchant K, ed. Laboratory Hematology Practice. Wiley Blackwell Publishing; 2012:435-446)

PDF Report

No

Day(s) Performed

Monday through Friday

Report Available

1 to 3 days

Specimen Retention Time

7 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

Fees & Codes

Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test has been modified from the manufacturer's instructions. Its performance characteristics were determined by

Mayo Clinic in a manner consistent with CLIA requirements. This test has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

85335

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
F9_IS	Factor IX Inhib Scrn	30086-3

Result ID	Test Result Name	Result LOINC® Value
7802	Factor IX Inhib Scrn	30086-3