



Test Definition: BART

Bartonella Antibody Panel, IgG and IgM, Serum

Overview

Useful For

Diagnosis of *Bartonella* infection, especially in the context of a cat scratch

Testing Algorithm

For information see [Infective Endocarditis: Diagnostic Testing for Identification of Microbiological Etiology](#).

Special Instructions

- [Infective Endocarditis: Diagnostic Testing for Identification of Microbiological Etiology](#)

Highlights

This assay can be used as an aid to diagnose recent or past infection with *Bartonella henselae* or *Bartonella quintana*.

Method Name

Immunofluorescence Assay (IFA)

NY State Available

Yes

Specimen

Specimen Type

Serum

Specimen Required

Supplies: Sarstedt Aliquot Tube 5 mL (T914)

Collection Container/Tube:

Preferred: Serum gel

Acceptable: Red top

Submission Container/Tube: Plastic vial

Specimen Volume: 0.5 mL

Collection Instructions: Centrifuge and aliquot serum into plastic vial.

Forms

If not ordering electronically, complete, print, and send [Infectious Disease Serology Test Request](#) (T916) with the specimen.

Specimen Minimum Volume

0.15 mL

Reject Due To

Gross hemolysis	Reject
Gross lipemia	Reject

Specimen Stability Information

Specimen Type	Temperature	Time	Special Container
Serum	Refrigerated (preferred)	30 days	
	Frozen	30 days	

Clinical & Interpretive

Clinical Information

Bartonella henselae and *Bartonella quintana* are small, rod-shaped, pleomorphic, gram-negative bacteria. The human body louse (*Pediculus humanis*) is the proposed vector for *B. quintana*. No animal reservoir has been determined for *B. quintana*. The domestic cat is believed to be both a reservoir and vector for *B. henselae*. Cats may infect humans directly through scratches, bites, or licks, or indirectly through an arthropod vector. Humans remain the only host in which *Bartonella* infection leads to significant disease.

The site of entry for *Bartonella* is through openings in the skin. Microscopically, *Bartonella* lesions appear as rounded aggregates that proliferate rapidly. These aggregates are masses of *Bartonella* bacteria. Warthin-Starry staining has shown that *Bartonella* organisms can be present within the vacuoles of endothelial cells, in macrophages, and between cells in areas of necrosis. Occasionally organisms are seen in the lumens of vessels. While cutaneous lesions are common, disseminated tissue infection by *Bartonella* has been seen in the blood, lymph nodes, spleen, liver, bone marrow, and heart. *B. henselae* has been associated with cat scratch disease (CSD), peliosis hepatitis (PH), bacillary angiomatosis (BA), and endocarditis. *B. quintana* has been associated with trench fever, BA, and endocarditis. BA is a vascular proliferative disease usually involving the skin and regional lymph nodes.

CSD begins as a cutaneous papule or pustule that usually develops within a week after animal contact. Regional lymphadenopathy follows and is the predominant clinical feature of CSD. Trench fever was a significant problem during World War I and World War II and is characterized by a relapsing fever and severe pain in the shins. PH and febrile bacteremia syndrome are both syndromes that have afflicted patients with AIDS and patients who are immunocompromised. While trench fever and CSD are usually self-limiting illnesses, the other *Bartonella*-associated diseases can be life-threatening.

Interest in *B. quintana* and *B. henselae* has recently increased since its increased prevalence in patients with AIDS, a transplanted organ, or suppressed immunity.

Reference Values

BARTONELLA HENSELAE

IgG: <1:128

IgM: <1:20

BARTONELLA QUINTANA

IgG: <1:128

IgM: <1:20

Interpretation

A positive immunofluorescence assay (IFA) IgM (titer >1:20) suggests a current infection with either *Bartonella henselae* or *Bartonella quintana*.

A positive IgG (titer >1:128) suggests a current or previous infection. Increases in IgG titers in serial specimens suggest active infection.

Normal serum specimens usually have an IgG titer of less than 1:128. However, 5% to 10% of healthy controls exhibit a *B. henselae* and *B. quintana* titer of 1:128. Sera from healthy volunteers rarely show titers of 1:256 or greater. IgM titers in normal serum are typically less than 1:20. IgM titers at 1:20 or greater have not been seen in the normal population.

Molecular testing of tissue for *Bartonella* species nucleic acid is recommended in cases of suspected endocarditis.

Clinical Reference

1. Rodino KG, Stone E, Saleh OA, Theel ES. The Brief case: Bartonella henselae endocarditis-a case of delayed diagnosis. J Clin Microbiol. 2019;57(9). e00114-19. doi:10.1128/JCM.00114-19
2. Wolf LA, Cherry NA, Maggi RG, Breitschwerdt EB. In pursuit of a stealth pathogen: Laboratory diagnosis of bartonellosis. Clin Micro News. 2014;36(5):33-39

Performance**Method Description**

The Euroimmun indirect immunofluorescence test is a standardized, in vitro assay for the determination of specific antibodies against *Bartonella henselae* and *Bartonella quintana*. BIOCHIP Mosaics are coated with *B. henselae* and *B. quintana* infected cells positioned next to each other in one reaction field. Samples are diluted and incubated on the substrate slides. If the reaction is positive, specific antibodies of class IgG and IgM attach to the antigens. In a second step, the attached antibodies are stained with fluorescein-labeled anti-human antibodies and made visible using fluorescence microscopy. Semiquantitative endpoint titers are obtained by testing serial dilutions of positive specimens. (Unpublished Mayo method)

PDF Report

No

Day(s) Performed

Monday through Saturday

Report Available

Same day/1 to 3 days

Specimen Retention Time

14 days

Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Superior Drive

Fees & Codes**Fees**

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

CPT Code Information

86611 x 4

LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
BART	Bartonella Ab Panel, IgG and IgM	90251-0

Result ID	Test Result Name	Result LOINC® Value
15659	Bart Henselae IgG	In Process
15660	Bart Henselae IgM	In Process
15661	Bart Quintana IgG	In Process
15662	Bart Quintana IgM	In Process