



## Test Definition: C5DCU

C5-DC Acylcarnitine, Quantitative, Random,  
Urine

### Overview

#### Useful For

Evaluation of patients with an abnormal newborn screen showing elevations of glutarylcarnitine

Diagnosis of glutaric aciduria type 1 deficiency

#### Genetics Test Information

Elevated glutarylcarnitine (C5-DC) in plasma or newborn screening blood spots is due to glutaric aciduria type 1 (GA-1), also known as glutaric acidemia type 1.

Urine C5-DC is a biochemical marker of GA-1 that appears to be elevated even in low excretors, who are affected patients with normal levels of glutaric acid in urine.

#### Method Name

Flow Injection Analysis-Tandem Mass Spectrometry (FIA-MS/MS)

#### NY State Available

Yes

### Specimen

#### Specimen Type

Urine

#### Ordering Guidance

This second-tier test is used specifically to evaluate a newborn screening elevation of glutarylcarnitine and **must not** be ordered with either C4U / C4 Acylcarnitine, Quantitative, Random, Urine or C5OHU / C5-OH Acylcarnitine, Quantitative, Random, Urine.

For general screening for metabolic disorders, see OAU / Organic Acids Screen, Random, Urine; ACRN / Acylcarnitines, Quantitative, Plasma; and AAQP / Amino Acids, Quantitative, Plasma.

#### Necessary Information

Patient's age, family history, clinical condition (asymptomatic or acute episode), diet, and drug therapy information is requested but not required.

#### Specimen Required

**Patient Preparation:** If clinically feasible, discontinue L-carnitine supplementation at least 72 hours before specimen collection.

**Supplies:** Urine Tubes, 10 mL (T068)

**Collection Container/Tube:** Clean, plastic urine collection container

**Submission Container/Tube:** Plastic, 10-mL urine tube

**Specimen Volume:** 5 mL

**Collection Instructions:**

1. Collect a random urine specimen.
2. Freeze specimen immediately.

**Forms**

[If not ordering electronically, complete, print, and send a Biochemical Genetics Test Request \(T798\)](#) with the specimen.

**Specimen Minimum Volume**

1 mL

**Reject Due To**

All specimens will be evaluated at Mayo Clinic Laboratories for test suitability.

**Specimen Stability Information**

Specimen Type	Temperature	Time	Special Container
Urine	Frozen (preferred)	7 days	
	Refrigerated	24 hours	

**Clinical & Interpretive**

**Clinical Information**

An isolated elevation of glutaryl-carnitine (C5-DC) in plasma or newborn screening blood spots is related to a diagnosis of glutaric aciduria type 1 (GA-1), also known as glutaric acidemia type 1. GA-1 is caused by a deficiency of glutaryl-CoA dehydrogenase. Diagnostic testing by acylcarnitine analysis, including the evaluation of C5-DC in urine, is helpful to determine if a patient has GA1. Urinary excretion of C5-DC is a specific biochemical marker of GA-1 that appears to be elevated even in low excretors, those patients who are affected but have normal levels of glutaric acid in urine.

Glutaric aciduria type 1 is an autosomal recessive disorder due to variants in the *GCDH* gene. It is characterized by bilateral striatal brain injury leading to dystonia, often a result of acute neurologic crises triggered by illness. Many affected individuals also have progressive macrocephaly. Dietary treatment and aggressive interventions during time of illness are recommended to try to prevent or minimize neurologic injury, which is most likely to occur in infancy and early childhood. Prevalence is approximately 1 in 100,000 individuals.

The American College of Medical Genetics and Genomics newborn screening work group published diagnostic algorithms for the follow-up of infants who had a positive newborn screening result. For more information, see the Practice Resources: ACT Sheets and Algorithms at [www.acmg.net](http://www.acmg.net).

**Reference Values**

---

<1.54 millimoles/mole creatinine

**Interpretation**

Elevated excretion of glutarylcarnitine is a specific biochemical marker of glutaric aciduria type 1 that is elevated in affected patients, apparently even in low excretors, ie, those affected individuals with normal levels of glutaric acid in urine.

**Cautions**

The results of urine acylcarnitines typically are not informative when the patient is receiving L-carnitine supplements.

**Clinical Reference**

1. Miller MJ, Cusmano-Ozog K, Oglesbee D, Young S. ACMG Laboratory Quality Assurance Committee: Laboratory analysis of acylcarnitines, 2020 update: a technical standard of the American College of Medical Genetics and Genomics (ACMG). *Genet Med.* 2021;23(2):249-258
2. Larson A, Goodman S. Glutaric Acidemia Type 1. In: Adam MP, Feldman J, Mirzaa GM, et al, eds. *GeneReviews* [Internet]. University of Washington, Seattle; 2019. Accessed December 16, 2025. Available at: <https://www.ncbi.nlm.nih.gov/books/NBK546575/>
3. Guenzel AJ, Hall P, Scott AI, et al. The low excretor phenotype of glutaric acidemia type I is a source of false negative newborn screening results and challenging diagnoses. *JIMD Rep.* 2021;60(1)67-74

**Performance****Method Description**

Acylcarnitines, including glutarylcarnitine, are determined in urine by flow injection analysis tandem mass spectrometry using acetyl-d3-carnitine, propionyl-d3-carnitine, butyryl-d3-carnitine, octanoyl-d3-carnitine, dodecanoyl-d3-carnitine, and palmitoyl-d3-carnitine as internal standards. The supernatant is evaporated and the residue treated with n-butanolic hydrochloric acid yielding the acylcarnitines for analysis as their n-butyl esters. (Tortorelli S, Hahn SH, Cowan TM, et al. The urinary excretion of glutarylcarnitine is an informative tool in the biochemical diagnosis of glutaric acidemia type I. *Mol Genet Metab.* 2005;84[2]:137-143; Miller MJ, Cusmano-Ozog K, Oglesbee D, Young S. ACMG Laboratory Quality Assurance Committee. Laboratory analysis of acylcarnitines, 2020 update: a technical standard of the American College of Medical Genetics and Genomics [ACMG]. *Genet Med.* 2021;23[2]:249-258)

**PDF Report**

No

**Day(s) Performed**

Monday, Wednesday, Friday

**Report Available**

2 to 5 days

**Specimen Retention Time**

1 month

## Performing Laboratory Location

Mayo Clinic Laboratories - Rochester Main Campus

## Fees & Codes

### Fees

- Authorized users can sign in to [Test Prices](#) for detailed fee information.
- Clients without access to Test Prices can contact [Customer Service](#) 24 hours a day, seven days a week.
- Prospective clients should contact their account representative. For assistance, contact [Customer Service](#).

### Test Classification

This test was developed and its performance characteristics determined by Mayo Clinic in a manner consistent with CLIA requirements. It has not been cleared or approved by the US Food and Drug Administration.

### CPT Code Information

82017

### LOINC® Information

Test ID	Test Order Name	Order LOINC® Value
C5DCU	C5-DC Acylcarnitine, QN, U	54279-5

Result ID	Test Result Name	Result LOINC® Value
88831	C5-DC Acylcarnitine, QN, U	54279-5
28126	C5-DC Interpretation	59462-2
34470	Reviewed By	18771-6